



Strawberry pre plant meetings, 2019

Pollination recommendations
Pest management considerations

Hannah Burrack, Professor & Extension Specialist

Jeremy Slone, PhD Graduate

Laura Kraft, PhD Student

Department of Entomology & Plant Pathology

Topics


- **Strawberry pollination**
 - Who are the Pollinators?
 - Farm Management practices
 - Pollinator Abundance and Diversity
 - Impacts on berry Production
- **Pest management considerations**
 - Spider mites
 - When is SWD a concern?



Screening spider mites for resistance




Screening spider mites for resistance



Mites collected from two strawberry fields, one watermelon field, and our laboratory population (4 populations total) appear highly susceptible to all ovicidal materials

Screening spider mites for resistance




***No different than control**

****Better than control, but weaker than other materials**

Greater variability among adulticides. Brigade and Vendex ineffective.

SWD in strawberries

- **Spotted wing drosophila (SWD)** is not a consistent pest in spring-fruiting strawberries
- **Fall fruiting berries** are at high risk
- In most years, the small infestation present can be managed through cultural control



Two types of studies


Field-Infested Fruit




Laboratory-Infested Fruit




Field-infested Strawberry




Field-infested Blueberry







Field-Infested Blackberry



Field-Infested Raspberry



Laboratory-Infested Strawberry




Topics

- **Strawberry pollination**
 - Who are the Pollinators?
 - Farm Management practices
 - Pollinator Abundance and Diversity
 - Impacts on berry Production
- **Pest management considerations**
 - Spider mites
 - When is SWD a concern?



Strawberry pollination




Poor Pollination



Why is pollinator diversity important?

What factors influence pollination?

pollinator behavior,
variable weather,
timing within the day/season, frost
protection




Who are the pollinators?

Managed Bees:

Honey Bees

- Hive nester
- Social species



Who are the pollinators?

Wild Bees:

- We collected 12 different bee genera.

<u>Agopostemon:</u>	15
<u>Andrena:</u>	91
<u>Augochlorella:</u>	42
<u>Halictus:</u>	63
<u>Lasioglossum:</u>	970
<u>Melissodes:</u>	92
<u>Megachilie:</u>	4
<u>Ceratina:</u>	3
<u>Osmia:</u>	8
<u>Calliopsis:</u>	1
<u>Epeolus:</u>	2
<u>Eucera:</u>	2

Who are the pollinators?

Wild Bees:

Lasioglossum spp.

- Ground nester
- Solitary species

Collected at least ~13 different *Lasioglossum* species/ species groups in 2017.



Sam Droege



Who are the pollinators?

Wild Bees:

Andrena spp.

Halictus spp.



Who are the pollinators?

Wild Bees:

Augochlorella
spp.



Sam Droege

Who are the pollinators?

Other insects:

Syrphid flies



Who are the pollinators?

Other insects:

Other flies?




Tom Murray



Judy Gallagher

Integrating pest and pollinator management




How does

**pesticide use, honey bee
stocking rate, & landscape**

**impact bee health and strawberry
pollination?**

Integrating pest and pollinator management



Pesticide Use

2017

- **13 conventional**
- **7 Organic**

- **6.4 average apps.**
 - 0 low - 24 high
 - 6 no spray sites

2018

- **12 conventional**
- **6 Organic**

- **5.8 average apps.**
 - 0 low - 21 high
 - 6 no spray sites

Pesticide Use

2017

- 13 conventional
- 7 Organic

- 6.4 average apps.
 - 0 low - 24 high
 - 6 no spray sites

- Herbicide avg – 0.32
- Insecticide avg – 0.89
- Acaricide avg – 0.89
- Fungicide avg – 4.3

2018

- 12 conventional
- 6 Organic

- 5.8 average apps.
 - 0 low - 21 high
 - 6 no spray sites

- Herbicide avg – 0.24
- Insecticide avg – 0.94
- Acaricide avg – 0.82
- Fungicide avg – 3.8

Pesticide Use

- *Conventional*
- *High intensity*
- *Organic*
- *Low intensity*

Landscape

- *Urban*
- *Forest*
- *Agricultural*
- *Natural*

Honey Bee Stocking Rate



500m & 1500m

Landscape

- Ag 500m – 1.6% to 78%
 - Ag 1500m – 1.3% to 78%
-
- Nat 500m – 2.8% to 54%
 - Nat 1500m – 3.1% to 75%

Pesticide Use

- *Conventional*
- *Organic*
- *High intensity*
- *Low intensity*

Landscape

- *Urban*
- *Agricultural*
- *Forest*
- *Natural*

Honey Bee Stocking Rate

- *Hives / acre*



Honey Bee Stocking Rate

2017

- 1.03 avg hives/acre
- 0 low to 4 high
- 9/20 with none


2018

- 0.76 avg hives/acre
- 0 low to 4 high
- 9/18 with none


Measuring pollinators



Pollinator abundance




Pollinator abundance




Pollinator Phenology

2017

Flies




Native Bees




Pollinator Phenology

2018


Flies



Native Bees



Pesticide intensity



- Bee abundance decreased with increasing pesticide application number.
 - -0.1858 ± 0.044
- Bee abundance was higher on conventional farms.
 - 1.33 ± 0.61

Landscape

- Bee abundance was lower when there was greater proportions of natural land at 1500m.

- -2.94 ± 1.39

Landscape composition may be less important than factors at a finer scale.

(floral diversity/abundance or available bare soil)



500m & 1500m

Honey bee stocking rate

- Honey bee density had a negative effect or no effect on native bee abundance captured via sweep net.


- -0.47 ± 0.18

- Honey bee density had a positive effect on native bee abundance captured via pan trap.

- 0.74 ± 0.16



Native bee diversity



2017

- **1.78 avg. score**
- **1 low to 3.35 high**

2018

- **1.71 avg score**
- **0 low to 3.46 high**

**Genus diversity did not vary
with farm management**

Who are the pollinators?

Wild bee abundance and diversity varies along the east coast, but community composition is somewhat similar.

- *NY (Finger Lakes region) – 14 farms, 2 years (organic/low intensity) – pan traps set out for 72hr, twice during the season.
 - <https://doi.org/10.1016/j.agee.2015.05.004>
- *NC Sampling – 18-20 farms, 2 years (Conv. & Org.) – pan traps set out for 6hr and sweep net sampling for 1.5 hours, three times during the season.

Wild bee genera	NY	NC
Agapostemon	2	15
Andrena	428	91
Augochlora	1	0
Augochlorella	85	42
Bombus	3	0
Calliopsis	0	1
Ceratina	37	3
Epeolus	0	2
Eucera	0	2
Halictus	26	63
Hoplitis	1	0
Hylaeus	5	0
Lasioglossum	364	970
Megachilie	0	4
Melissodes	0	92
Nomada	38	0
Osmia	9	8
Sphecodes	3	0
Xylocopa	1	0

Non-bee Pollinators



Pesticide intensity



- Pesticide application number had nearly no effect on fly abundance or diversity.
- Conventional management had a negative effect on abundance, but a positive effect on 2018 diversity
 - **Abundance:** -0.45 ± 0.19
 - **Diversity:** 2.0 ± 0.67



Landscape



- Greater natural land cover had a positive effect on abundance, but a negative effect on 2018 diversity.

- Abundance: 0.58 ± 0.29

- Diversity: -3.18 ± 1.33

- Greater agricultural land cover had a positive effect on 2018 diversity.

- 2.13 ± 0.89



500m & 1500m

Honey bee stocking rate



- **Greater honey bee density had a positive effect on fly abundance, but a negative effect on 2017 diversity.**

- **Abundance : 0.20 ± 0.07**
- **Diversity: -0.63 ± 0.17**

Strawberry size and quality





Strawberry size and quality



- **Neither native bee abundance nor diversity impacted weight or symmetry...**
 - **Reaching complete pollination?**



Strawberry size and quality

2017





Strawberry size and quality

2017





Strawberry size and quality

2018



Strawberry size and quality


2018



Strawberry size and quality



- Higher honey bee density had a **negative** effect on strawberry weight.



Strawberry size and quality

- Greater proportions of agricultural land had a **positive** effect on berry symmetry



Pollination Conclusions

- **Pesticide (fungicide) use can negatively impact pollinators in strawberries**
- **Insect pollinators do not appear to benefit strawberry weight or symmetry**
- **Therefore pesticide impacts on pollinators may be more important for other crops within a farm.**
- **Stocking too many honey bees may decrease strawberry weight**
- **Unless honey bees are needed for other crops at the same time, stocking them may be unnecessary in strawberries**

*Read us @ entomology.ces.ncsu.edu
Like us @ facebook.com/NCsmallFruitIPM
Follow us @NCsmallFruitIPM*

